

Prevalence of elevated blood pressure in children and adolescents in Africa: a systematic review and meta-analysis

Jean Jacques Noubiap, Mickael Essouma, Jean Joel Bigna, Ahmadou M Jingi, Leopold N Aminde, Jobert Richie Nansseu



Summary

Background Despite substantial attention paid to the threat of elevated blood pressure in children and adolescents in high-income countries and the epidemic of hypertension in African adult populations, data on the burden of elevated blood pressure in African children and adolescents have not yet been synthesised. We did a systematic review and meta-analysis to provide estimates of the prevalence of elevated blood pressure and assess associated factors among children and adolescents in Africa.

Methods We searched Embase, PubMed, African Journals Online, and African Index Medicus to identify articles published from Jan 1, 1996, to Feb 2, 2017, and searched the reference list of retrieved articles. Each study was independently reviewed for methodological quality. We used a random-effects model to estimate the prevalence of elevated blood pressure across studies and heterogeneity (I^2) was assessed via the χ^2 test on Cochran's Q statistic. This review is registered with PROSPERO, number CRD42015019029.

Findings We included 51 studies in qualitative synthesis and 25 in the meta-analysis reporting data of a pooled sample of 54 196 participants aged 2–19 years. Study quality was high with only four medium-quality studies and no low-quality studies. Prevalence of elevated blood pressure varied widely across studies (range 0·2–24·8%). The pooled prevalence of elevated blood pressure (systolic or diastolic blood pressure ≥ 95 th percentile) was 5·5% (95% CI 4·2–6·9), whereas that of slightly elevated blood pressure (systolic or diastolic blood pressure ≥ 90 th percentile and < 95 th percentile) was 12·7% (2·1–30·4). The prevalence of elevated blood pressure was largely associated with body-mass index (BMI), with a prevalence of elevated blood pressure six times higher in obese (30·8%, 95% CI 20·1–42·6) versus normal-weight children (5·5%, 3·1–8·4; $p < 0\cdot0001$).

Interpretation This study suggests a high prevalence of elevated blood pressure among children and adolescents in Africa, with overweight and obesity being an important risk factor. Efforts to address this burden of elevated blood pressure in children and adolescents should mainly focus on primary prevention at the community level, by promoting healthy lifestyles and avoiding other cardiovascular risk factors, especially overweight and obesity. This study also stresses the need for more elaborate studies using uniform and reliable diagnostic methods to reliably map the burden of elevated blood pressure in children and adolescents in Africa.

Funding None.

Copyright © The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

Hypertension is the greatest modifiable risk factor for cardiovascular disease, disability, and deaths worldwide. Globally, it affects about 1 billion adults and is associated with more than 9 million deaths annually.¹ Furthermore, hypertension was the largest contributor to global disability-adjusted life-years (DALYs) in 2015, accounting for 9·2% (95% uncertainty interval 8·3–10·2) of DALYs for men and 7·8% (6·9–8·7) of DALYs for women.² Once uncommon in African populations, the burden of hypertension has been continuously rising over the past decades in these populations, driven by reduced physical activity, unhealthy diet, and obesity.³ According to WHO, Africa has the highest age-standardised prevalence of hypertension, with 46% of adults older than 25 years being affected.¹

Although hypertension is less common in children than in adults, its burden has increased in children and

adolescents in some high-income countries over the past decade, mainly due to the surge in obesity prevalence.^{4–10} Indeed, reports from epidemiological studies on blood pressure in children and adolescents done during the past decade have shown a substantial increase in blood pressure levels and in the prevalence of hypertension in these age groups, with rates ranging from 1% to 5% in some western countries.^{4–10} However, these increasing trends have not been observed in all countries studied.¹¹

Hypertension in children and adolescents is becoming a major concern, not only because of its rising prevalence, but also because of evidence suggesting that hypertension tracks from childhood to adulthood. Almost half of adults with hypertension had elevated blood pressure values during childhood.^{8–13} Furthermore, several studies have suggested that elevated blood pressure in childhood correlates with carotid intima-media thickness, atherosclerosis,

Lancet Public Health 2017; 2: e375–86

See [Comment](#) page e346

Department of Medicine, University of Cape Town and Groote Schuur Hospital, Cape Town, South Africa (J J Noubiap MD); Department of Internal Medicine and Specialties (M Essouma MD, A M Jingi MD) and Department of Public Health (J R Nansseu MD), Faculty of Medicine and Biomedical Sciences, University of Yaoundé I, Yaoundé, Cameroon; Department of Epidemiology and Public Health, Centre Pasteur of Cameroon, Yaoundé, Cameroon (J J Bigna MD); Faculty of Medicine, University of Paris Sud XI, Le Kremlin Bicêtre, France (J J Bigna); and School of Public Health, Faculty of Medicine, The University of Queensland, Brisbane, QLD, Australia (L N Aminde MD)

Correspondence to: Dr Jean Jacques Noubiap, Department of Medicine, University of Cape Town and Groote Schuur Hospital, Cape Town 7925, South Africa noubiapjj@yahoo.fr

Research in context

Evidence before this study

There is evidence that hypertension and its cardiovascular complications track from childhood to adulthood. Little is known about the burden of hypertension in children and adolescents in Africa. Only one previous meta-analysis at the global level has provided an estimate of the prevalence of hypertension in adolescents in Africa. Unfortunately, this review included only three studies from Africa and the derived prevalence of 25.5% was therefore questionable.

Added value of this study

To the best of our knowledge, this study is the first comprehensive summary of available high quality surveys estimating the point prevalence of elevated blood pressure in children and adolescents in Africa. Our study shows that elevated blood pressure is not infrequent in these populations, with overweight and obesity being a driver of the condition. The weak influence of the gross domestic product per capita on the prevalence of elevated blood pressure suggests that elevated blood pressure is equally affecting children and adolescents in poorer and less poor countries in Africa.

The influence of socioeconomic level on the occurrence of elevated blood pressure during childhood and adolescence might be more apparent at the individual level than at the country level. Another salient finding is the increased prevalence of elevated blood pressure in children and adolescents in rural areas, which deserves further investigation.

Implications of all the available evidence

We hope results from this research will prompt implementation of appropriate policy towards improving awareness, diagnosis, and proper management of hypertension in African children and adolescents. In the absence of sufficient evidence to recommend universal blood pressure screening, we advocate for targeted identification of elevated blood pressure according to the clinical context, and early treatment of cases with secondary hypertension. At the population level, efforts should focus on primary prevention of elevated blood pressure by promoting weight control, healthy diet, and regular physical activity. Strategies for blood pressure control and cardiovascular disease prevention in childhood and adolescence might be integrated in existing programmes.

left ventricular hypertrophy, and kidney failure in adulthood.^{14–21} Consequently, early diagnosis and control of hypertension in childhood are likely to have an important effect on long-term outcomes of hypertension-related cardiovascular complications.

Although epidemiological data on hypertension and cardiovascular disease in adults in Africa are growing, the prevalence of elevated blood pressure and its drivers in children and adolescents in Africa remain largely unknown. Hence, we did a systematic review of the literature and a meta-analysis to provide a reliable estimate of the prevalence of elevated blood pressure and identify associated factors among children and adolescents in Africa.

Methods

Search strategy and selection criteria

For this systematic review and meta-analysis, JJN, ME, and LNA, with the help of an expert librarian, did a comprehensive search of PubMed, Embase, African Journals Online, and African Index Medicus to identify all relevant articles on elevated blood pressure in children and adolescents in Africa published in English and French from Jan 1, 1996, to Feb 2, 2017. We focused on the past 20 years to provide recent estimates and to reduce the heterogeneity between studies. We conceived and applied a search strategy based on the combination of relevant terms and names of each of the 54 African countries and African subregions such as “north Africa” or “central Africa”. Names of African countries were included in English and in the official languages of respective countries (eg, “Ivory Coast” and

“Côte d’Ivoire”). For countries whose names have changed over time, both the old and the new name were used (eg, “Zaire” and “Democratic Republic of the Congo”). Terms used for hypertension included the following: “hypertension”, “blood pressure”, “systolic hypertension”, and “diastolic hypertension”. For children and adolescents, we used “child”, “children”, “childhood”, “adolescent”, “teens”, “teenage”, “youth”, “infant”, and “pediatric”. The main search strategy done in PubMed is available in the review protocol.²² This search strategy was adapted to fit with other databases. To supplement these bibliographic database searches, references of all relevant research articles and reviews were also scrutinised to identify additional potential data sources.

To be included in this systematic review, primary studies had to be observational studies of children and adolescents aged 1–19 years residing in African countries, irrespective of ethnic, socioeconomic, and educational backgrounds, reporting the prevalence of elevated blood pressure or with enough data to compute these estimates. We excluded studies on non-systemic hypertension (intracranial or pulmonary hypertension), those done among populations of African origin residing outside Africa, studies in subgroups of participants selected on the basis of presence of hypertension (eg, clinical trials or case-control studies), and studies including both adult and paediatric populations in which it was not possible to disaggregate data for children or adolescents. We also excluded case series with a small sample size (<50 participants), letters, reviews, commentaries, editorials, and studies without primary data or explicit description of methods. For studies published in more

than one report (duplicates), we considered the most comprehensive report with the largest sample size.

Two investigators (JJN and ME) independently screened the titles and abstracts of articles retrieved from the literature search, and full texts of articles found potentially eligible were obtained and further assessed for final inclusion. All duplicates were removed during the study selection process. Disagreements were resolved through discussions between investigators until a consensus was reached.

Data extraction

Two investigators (JJN and ME) independently extracted relevant data from individual studies using a preconceived and standardised data extraction form.

Information extracted included first author's name, year of publication, recruitment period, area (rural vs urban), country, study design, setting, sample size, mean or median age, age range, proportion of male participants, proportion of participants with obesity, ascertainment of elevated blood pressure and diagnostic criteria, and the number of participants with slightly elevated blood pressure (systolic or diastolic blood pressure ≥ 90 th percentile and < 95 th percentile) and elevated blood pressure (systolic or diastolic blood pressure ≥ 95 th percentile). We assigned a United Nations Statistics Division (UNSD) African region (Central [refers to Central Africa], Eastern, Northern, Southern, and Western) to each study regarding the country of recruitment. We retrieved the gross domestic product per capita (GDP) in US dollars of each study from the World Bank database and corresponded it to the GDP of the year of recruitment of each study's country. For studies with recruitment in more than one year, we considered the median GDP of the corresponding country in the years of recruitment. Disagreements between authors were reconciled through discussion and consensus. Where relevant data were not available, we directly contacted the corresponding author to request the information.

Data analysis

We used a meta-analysis to summarise prevalence data. We determined SEs for study-specific estimates from the point estimate and the appropriate denominators. We pooled the study-specific estimates using a random-effects meta-analysis model to obtain an overall summary estimate of the prevalence across studies,²³ after stabilising the variance of individual studies with the use of the Freeman-Tukey double arcsine transformation.²⁴

We evaluated the methodological quality of included studies using the tool developed by Hoy and colleagues.²⁵ We assigned each item a score of 1 (yes) or 0 (no), and summed scores across items to generate an overall quality score that ranged from 0 to 10. According to the overall scores, we classified studies as having a low (> 8), moderate (6–8), or high (≤ 5) risk of bias. Two investigators (JJN and JJB) independently assessed study quality, with disagreements resolved by consensus.

We assessed heterogeneity using the χ^2 test on Cochran's Q statistic²⁶ and quantified heterogeneity by calculating the I^2 (with values of 25%, 50%, and 75% representing low, medium, and high heterogeneity, respectively).²⁷ To be included in the meta-analysis, studies had to define elevated blood pressure (or hypertension) as systolic or diastolic blood pressure greater than or equal to the 95th percentile (for all ages) or slightly elevated blood pressure (or pre-hypertension) as systolic or diastolic blood pressure in the 95th percentile but greater than or equal to the 90th percentile, with a random selection of participants with no specific disease or profile, low risk of bias in their methodological quality, and prospective data collection. If substantial heterogeneity was detected, we did subgroup analysis when possible to investigate the possible sources of heterogeneity using the following grouping variables: age group, sex, study setting, UNSD African region, number of blood pressure measurements, representativeness of sample, sample size, period of participants' recruitment, and body-mass index (BMI). Subgroup comparisons used the Q test based on ANOVA. We considered a subgroup difference p value less than 0.05 to be indicative of significant difference between subgroups. We used meta-regression to measure the association between prevalence of elevated blood pressure and GDP. We adjusted for BMI, considering the strong association between GDP and obesity. In the case of non-normal distribution of GDP, we planned a $\log_{10}$ -transformation of the GDP for meta-regression and presentation in a scatter plot. We assessed the presence of publication bias using funnel plots and the formal Egger test. We considered an Egger test p value less than 0.10 to be indicative of significant publication bias.²⁸ We assessed inter-rater agreement for study inclusion and data extraction using Cohen's κ coefficient.²⁹ We analysed data using Stata version 13.0 for Windows. The meta-analysis was preceded by a qualitative synthesis of data.

This review was registered in the PROSPERO International Prospective Register of systematic reviews, registration number CRD42015019029 and its protocol has been published.²²

Role of the funding source

There was no funding source for this study. The corresponding author had full access to all study data and had final responsibility for the decision to submit the paper for publication.

Results

We identified 5891 records; after elimination of duplicates, 3543 records remained (figure 1). After screening titles and abstracts, we found 3468 records to be irrelevant and excluded them. The inter-rater agreement for study selection was high ($\kappa = 0.88$). We assessed full texts of the remaining 75 papers for eligibility, of which 28 were excluded. We included a total of 51 studies^{30–76} from 47 full-text articles in the review,

For the breakdown of African regions see <https://unstats.un.org/unsd/methodology/m49/>




Figure 1: Study selection

with only 25 of the studies^{30–50} considered in the pooled analysis. Inter-rater agreements were high for both study inclusion ($\kappa=0.93$) and data extraction (0.89).

The methodological quality and characteristics of included studies are presented in table 1. Three studies provided cross-sectional data of a cohort study and the other data were pulled out from cross-sectional studies. Data were from 13 countries distributed across the five UNSD regions of Africa. The definitions of elevated blood pressure varied widely across studies. With respect to our criteria, 40 studies were classified as having an acceptable definition for elevated blood pressure^{30–53,55–60,62,64,66–68,73,74} whereas ten studies used a non-acceptable definition^{54,61,63,65,69,70,72,75,76} and one did not report the definition used.⁷¹ Among these ten studies with non-acceptable definitions of elevated blood pressure, four studies defined elevated blood pressure as blood pressure of 120/80 mm Hg or higher,^{63,65,67,72} three as blood pressure greater than the mean plus two SDs,^{61,69,70} two as systolic or diastolic blood pressure within the 95th percentile for individuals younger than 18 years and blood pressure of 140/90 mm Hg or higher for individuals aged 18 years or older,⁷⁶ and one as blood pressure of 125/84 mm Hg or higher.⁷⁵ All 25 studies included in the meta-analysis had low risk of bias in their methodology, randomly selected participants, and were population based.^{30–50}

From the 51 studies included in the review, the prevalence of elevated blood pressure varied widely from 0.2% to 24.8%. Prevalence in different UNSD regions in Africa was 9.8–10.1% (Central), 6.5–17.1% (Eastern), 4.8–19.7% (Northern), 4.5–21.2% (Southern), and 0.2–24.8% (Western). Prevalence was between 4.1% and 11.2% in rural areas, 0.2% and 24.8% in urban areas,

and 2.6% and 21.2% in studies done in both areas. Among females, prevalence varied from 0.4% to 20.9% and among males from 0.6% to 25.0%. Prevalence widely varied by age group: 2–5 years (0.5–22.4%), 6–12 years (1.5–24.4%), and 13–19 years (1.8–21.2%). Prevalence of elevated blood pressure in the two studies with specific profile (overweight and obese) was high, at 17.4% and 19.7%. Prevalence of elevated systolic blood pressure ranged from 0.1% to 14.2%, and from 0.1% to 14.6% for elevated diastolic blood pressure. Prevalence of slightly elevated blood pressure retrieved from ten studies done in Nigeria, South Africa, Tunisia, and Congo (Brazzaville) varied from 1.4% to 39.1%.

Table 2 summarises results from meta-analyses. The overall prevalence of elevated blood pressure was 5.5% (95% CI 4.2–6.9) in a pooled sample of 54196 children and adolescents from 25 studies (figure 2). The funnel plot did not suggest any publication bias (appendix p 7) and this result is confirmed by the Egger test (table 2). The prevalence of slightly elevated blood pressure was about twice as high as overall prevalence (table 2, appendix p 10). The prevalence of elevated systolic blood pressure and that of elevated diastolic blood pressure were similar (table 2, appendix pp 8–9).

The GDP per capita was not normally distributed and therefore $\log_{10}$ -transformed (appendix p 11). Prevalence of elevated blood pressure generally increased slightly with the $\log_{10}$ -transformed GDP. This was confirmed by the results of meta-regression (coefficient 0.04, 95% CI 0.01–0.07; $p=0.015$; 25 studies). The $\log_{10}$ -transformed GDP explained only 0.13% of the variation in the prevalence of elevated blood pressure. After adjustment for mean BMI, the association remained significant (adjusted coefficient 0.04, 0.01–0.07; $p=0.017$; 18 studies), explaining only 0.16% of the variation in the prevalence of elevated blood pressure.

Publication bias was found for the following subgroup analyses: elevated diastolic blood pressure, setting, national representative studies, studies using mercury devices for blood pressure measurement, and studies with sample size higher than the median (table 2). Substantial heterogeneity was present within all subgroups (table 2).

Data for the UNSD region of Western Africa came from Nigeria (ten studies) and Côte d'Ivoire (one study). For Northern Africa, data came only from Tunisia (four studies). For the region of Southern Africa, data came only from South Africa (three studies). For Eastern Africa, data came from Seychelles (five studies) and Uganda (one study); and for Central Africa, data came only from Congo (Brazzaville; one study). The prevalence of elevated blood pressure in Western Africa was lower than that in other UNSD regions of Africa (table 2, figure 2). 16 studies presented data from urban areas and four from rural areas; the prevalence of elevated blood pressure was higher in rural areas (table 2; appendix p 12).

Prevalence of elevated blood pressure did not differ between age groups and sex (table 2; appendix pp 13–14).

See Online for appendix

The prevalence of elevated blood pressure was higher when blood pressure measurements were taken with a digital device compared with aneroid or mercury devices (table 2; appendix p 16). We also found a difference for measurement method: the prevalence with one reading was lower than that obtained with an average of two or three readings (table 2; appendix p 17).

With regard to BMI, the prevalence of elevated blood pressure among obese and overweight children was higher than among normal-weight and underweight populations (table 2, figure 3).

Discussion

This systematic review and meta-analysis of data from 25 studies involving 54 196 individuals found a pooled prevalence of elevated blood pressure of 5.5% and a pooled prevalence of slightly elevated blood pressure of 12.7% in children and adolescents in Africa. Increased BMI was largely associated with the prevalence of elevated blood pressure, with prevalence six times higher in obese children and adolescents than in those of normal weight.

The prevalence of elevated blood pressure found in this review is below the pooled prevalence estimate (25.5%, 95% CI 10.1–39.1) of elevated blood pressure in African adolescents found by a previous systematic review.⁷⁷ This discrepancy might be explained by difference in the number of surveys included in the pooled analyses (25 in the current study *vs* three in the other) as well as different age bands (2–19 years *vs* 10–19 years).⁷⁷ de Moraes and colleagues' systematic review⁷⁷ was not designed to estimate the prevalence of high blood pressure in adolescents in Africa specifically, but was a worldwide study to compare prevalence estimates of elevated blood pressure in adolescents across different regions. In their review,⁷⁷ Africa and Oceania (which had the most limited numbers of included studies) had the highest prevalence estimates when compared with other regions (North America, Latin America, Asia, and the Middle East; $p=0.001$). Nevertheless, the prevalence of elevated blood pressure found in our review is similar to the 1–5% prevalence reported in children and adolescents from high-income countries and other low-income and middle-income countries.^{4–10}

We found no difference between boys and girls in terms of prevalence of elevated blood pressure. This finding supplements results from the most recent meta-analysis⁷⁸ on hypertension in adults in Africa. Both studies suggest that male sex might not be a risk factor for elevated blood pressure in Africa, unlike reports from some developed countries.⁷⁹

A key finding of this study is the substantial association of the prevalence of elevated blood pressure with increased BMI. A number of epidemiological studies from Africa and other regions have previously reported this link between elevated blood pressure and increased BMI (obesity and overweight) both in adults

	All studies (n=51)	Studies included in the meta-analysis (n=25)
Methodological quality		
Timing of data collection	Prospective (n=50), retrospective (n=1)	All prospective
Risk of bias	Low risk (n=47), moderate risk (n=4)	All low risk
Selection of participants	Random selection (n=40), non-random (n=11)	All random
Representativeness	National (n=8), subnational (n=43)	National (n=7), subnational (n=18)
Population based	All	All
Characteristics		
Year of publication	1998–2016	2001–2016
Patient recruitment period	1995–2013	1997–2013
Countries	Algeria (n=2), Angola (n=1), Congo (Brazzaville) (n=1), Côte d'Ivoire (n=1), Democratic Republic of the Congo (n=1), Egypt (n=1), Morocco (n=1), Nigeria (n=24), Senegal (n=1), Seychelles (n=6), South Africa (n=6), Tunisia (n=5), and Uganda (n=1)	Congo (Brazzaville) (n=1), Côte d'Ivoire (n=1), Nigeria (n=10), Seychelles (n=5), South Africa (n=3), Tunisia (n=4), and Uganda (n=1)
UNSD African regions	Western (n=26), Northern (n=9), Eastern (n=7), Southern (n=7), and Central (n=2)	Western (n=11), Eastern (n=6), Northern (n=4), Southern (n=3), and Central (n=1)
Settings	Rural (n=6), urban (31), and both (n=14)	Rural (n=2), urban (n=14), and both (n=9)
Blood pressure values considered	One reading (n=2), lowest diastolic blood pressure and its matching systolic blood pressure after three measurements (n=1), average of two readings (n=20), and average of three readings (n=22); missing data (n=6)	One reading (n=1), average of two readings (n=8), and average of three readings (n=13); missing data (n=3)
Types of device for blood pressure measurement	Aneroid (n=7), digital (n=14), and mercury (n=24); missing data (n=6)	Aneroid (n=5), digital (n=3), and mercury (n=14); missing data (n=3)
Elevated blood pressure definition	Good definition* (n=40) and others (n=11)	Good definition* (n=25)
Mean age (years)	3.7–17.7 (from 45 studies)	3.7–16.9 (from 22 studies)
Age range (years)	1–19	2–19
Female participants	37.7–73.7% (from 44 studies†)	45.2–61.9%
Male participants	26.3–62.3% (from 44 studies†)	38.1–54.8%
Participants' specific profile or disease	Overweight (n=1), overweight and obese (n=1), and none (n=49)	None (n=25)
Gross domestic product per capita, median (IQR)	US\$2939.69 (2281.06–4038.07; from 41 studies)	US\$2829.05 (2310.86–5814.33)
UNSD=United Nations Statistics Division. *Systolic or diastolic blood pressure $\geq$ 95th percentile. †One study included only males.		

Table 1: Characteristics of included studies

and in children and adolescents.^{2,8,9,54} Although we could not accurately determine whether the link observed was causal or not, obesity and overweight are known risk factors for elevated blood pressure and high blood pressure-specific cardiovascular disease as well as all-cause cardiovascular disease in all age groups.^{8,9} Accordingly, increased BMI is probably an important risk factor for elevated blood pressure in children and adolescents in Africa. Notably, obese individuals have excess fat leading to elevated blood pressure via an increase in sympathetic activity and




Figure 2: Prevalence of elevated blood pressure in children and adolescents by UNSD African regions
UNSD=United Nations Statistics Division.

subsequent sodium reabsorption and increased peripheral vascular resistance.

GDP is generally associated with obesity,^{80–82} and obesity is strongly associated with elevated blood pressure.^{2,8,9,54} We found a weak influence of GDP on prevalence of elevated blood pressure in this study, suggesting that elevated blood pressure is equally affecting children and adolescents in poorer and less poor countries in Africa. This result has implications in terms of the coping capacities of countries; less developed countries can be disproportionately affected by the burden of disease, but tackling the burden of elevated blood pressure in children and adolescents in such countries in Africa might not require supplemental efforts and resources.

Prevalence of elevated blood pressure was higher in rural areas than in urban areas. This pattern of elevated blood pressure has also been reported in children and adolescents from Poland.⁸³ Environmental variations of blood pressure levels are difficult to explain. Nevertheless, two potential underlying mechanisms have been described: low socioeconomic status and obesity.^{77,83–86} Elevated blood pressure has been associated with low socioeconomic status in previous studies^{84–87} from low-income and middle-income countries including African countries, in both adult and paediatric populations. Along the same lines, a study⁸⁸ in South Africa has shown that improvement in socioeconomic status between infancy and adolescence has a protective effect on systolic

blood pressure in young adulthood. Rural area inhabitants often live in impoverished conditions with low household status, exposing them to increased stress perception, a factor that is known to enhance activation of the autonomic nervous system and subsequently

increase blood pressure. Lower socioeconomic conditions in rural areas might therefore account for the increased prevalence of elevated blood pressure in children and adolescents living in these settings, as compared to those in urban areas. In addition, studies^{89,90} in several

	Number of studies	Number of participants	Number of cases	Prevalence (95% CI)	I ²	p values		
						Heterogeneity	Subgroup difference	Egger test
Global analysis								
Overall elevated blood pressure	25	54 196	3569	5.5% (4.2–6.9)	97.9%	<0.0001	..	0.265
Elevated systolic blood pressure	10	18 473	938	3.6% (1.7–6.4)	98.6%	<0.0001	..	0.113
Elevated diastolic blood pressure	10	18 473	1395	3.0% (0.3–8.5)	97.7%	<0.0001	..	0.034
Slightly elevated blood pressure	5	8437	1746	12.7% (2.1–30.4)	99.6%	<0.0001	..	0.402
Subgroup analysis								
United Nations Africa region								
Overall	25	51 986	4069	5.5% (4.2–6.9)	97.9%	<0.0001	<0.0001	0.265
Central	1	603	61	10.1% (8.0–12.8)	..	..	..	Not estimable*
Eastern	6	21 108	2339	9.0% (7.7–10.5)	93.6%	<0.0001	..	0.071
Northern	4	7009	466	7.3% (4.7–10.5)	95.4%	<0.0001	..	0.181
Southern	3	3131	261	8.1% (4.3–13.1)	93.9%	<0.0001	..	0.915
Western	11	15 345	442	2.5% (1.4–3.8)	95.5%	<0.0001	..	0.910
Setting								
Both urban and rural	20	23 301	1196	5.4% (3.7–7.4)	97.4%	<0.0001	0.013	0.057
Urban	16	19 089	872	4.6% (2.8–6.7)	97.5%	<0.0001	..	0.173
Rural	4	4212	324	9.2% (6.2–12.8)	91.7%	<0.0001	..	0.105
Age group								
Overall	16	17 844	2013	8.7% (5.8–12.0)	98.0%	<0.0001	0.631	0.797
2–5 years	2	1656	233	10.7% (9.3–12.2)	99.9%	<0.0001	..	Not estimable*
6–12 years	4	2634	319	7.0% (0.4–20.0)	98.8%	<0.0001	..	0.752
13–19 years	10	13 554	1461	9.5% (6.8–12.4)	96.2%	<0.0001	..	0.466
Sex								
Overall	30	43 345	3418	6.6% (5.3–7.9)	96.7%	<0.0001	0.856	0.896
Male	15	21 065	1689	6.4% (4.6–8.5)	96.9%	<0.0001	..	0.246
Female	15	22 280	1729	7.0% (5.2–9.0)	96.8%	<0.0001	..	0.581
Sample representativeness								
Overall	25	51 986	4069	5.5% (4.2–6.9)	97.9%	<0.0001	0.065	0.532
National	7	32 216	2474	7.2% (6.0–8.4)	94.4%	<0.0001	..	0.074
Subnational	18	21 980	1095	4.9% (3.2–7.0)	97.7%	<0.0001	..	0.157
Device type for blood pressure measurement								
Overall	22	50 334	3478	5.9% (4.5–7.3)	97.7%	<0.0001	<0.0001	0.999
Mercury	14	39 401	2697	4.8% (3.6–6.2)	97.0%	<0.0001	..	<0.0001
Aneroid	5	7932	450	5.8% (2.0–11.5)	98.7%	<0.0001	..	0.325
Digital	3	2901	331	12.3% (8.6–16.6)	90.1%	<0.0001	..	0.259
Number of measurements								
Overall	22	48 028	3860	5.4% (4.2–6.8)	97.9%	<0.0001	<0.0001	0.144
One	1	1638	3	0.2% (0.1–0.5)	..	..	..	Not estimable*
Average of two	8	11 167	575	4.9% (3.1–7.2)	95.9%	<0.0001	..	0.092
Average of three	13	38 059	2822	6.4% (5.1–7.9)	96.7%	<0.0001	..	0.392

(Table 2 continues on next page)

(Table 2 continues on next page)

	Number of studies	Number of participants	Number of cases	Prevalence (95% CI)	I ²	p values		
						Heterogeneity	Subgroup difference	Egger test
(Continued from previous page)								
BMI†								
Overall	14	6829	469	14.4% (10.1–19.3)	95.4%	<0.0001	<0.0001	<0.0001
Underweight	1	219	16	7.3% (4.5–11.5)	..	..	..	Not estimable*
Normal	4	5091	224	5.5% (3.1–8.4)	92.6%	<0.0001	..	0.326
Overweight	5	1271	162	16.6% (9.1–25.9)	92.8%	<0.0001	..	0.119
Obese	4	248	67	30.8% (20.1–42.6)	64.9%	0.0359	..	0.010
Recruitment period								
Overall	18	47 199	3320	6.5% (5.2–8.0)	97.3%	<0.0001	0.054	0.302
1997–2006	8	30 114	2385	8.1% (6.8–9.4)	94.0%	<0.0001	..	0.619
2007–16	10	17 085	935	5.3% (3.3–7.8)	97.6%	<0.0001	..	0.844
Median sample size								
Overall	25	51 986	4069	5.5% (4.2–6.9)	97.9%	<0.0001	0.557	1.0
≤1569	13	11 708	735	5.9% (3.6–8.7)	97.1%	<0.0001	..	0.425
>1569	12	42 488	2834	5.1% (3.5–6.9)	98.4%	<0.0001	..	<0.0001

..=not applicable. BMI=body-mass index. *Because of insufficient observations. †Underweight: BMI less than 2 SDs below the WHO growth standard mean for age and sex; normal: BMI between 2 SDs below and 1 SD above the mean; overweight: BMI more than 1 SD above the mean; and obesity: BMI more than 2 SDs above the mean.

Table 2: Prevalence of elevated blood pressure among children and adolescents in Africa using random-effects model and double arcsine transformation

developed countries have shown that overweight and obesity are more prevalent in children and adolescents living in rural areas, representing a plausible explanation for the higher prevalence of elevated blood pressure in rural areas than in urban areas in these age groups. However, data on urban–rural differences in terms of prevalence of obesity in African countries are insufficient to support this assumption.

Taken collectively, elevated blood pressure is a health problem that might be largely driven by high BMI in children and adolescents in Africa. Some questions therefore arise from our study. First, can primary prevention of hypertension be beneficial for children and adolescents in Africa? Second, if necessary, what are the most suitable preventive strategies? Third, is treatment of essential hypertension necessary and for how long is it recommended?

Primary prevention of hypertension in children and adolescents aims to reduce the risk of cardiovascular disease and prevent hypertension in adulthood. Although high blood pressure is associated with detrimental cardiovascular effects in children and teens, no direct evidence exists to estimate the absolute risk of cardiovascular disease associated to a particular level of blood pressure in childhood. The robustness of hypertension tracking should be further evaluated given the difficulty of obtaining perfect blood pressure measurements, especially in children and adolescents.^{8–13} As a result, existing guidelines do not endorse primary prevention of hypertension in childhood through

systematic screening.⁸⁵ Nevertheless, early identification of children and adolescents with elevated blood pressure could help to reduce the hypertension epidemic in adults and cardiovascular disease across the continent. Primary prevention of elevated blood pressure in children and adolescents should first and foremost rely on population interventions addressing modifiable blood pressure determinants—mainly weight control, modification of unhealthy diet and promotion of regular physical activity. Blood pressure control and cardiovascular prevention programmes addressing modifiable risk factors are already in place in most African countries, but are mostly designated for adults. Their implementation should be extended to younger age groups. Existing programmes, such as child and maternal health programmes, and school programmes could be useful. Second, depending on resources available, primary prevention of elevated blood pressure at the individual level should rely on weight control (as weight excess is by far the main determinant of elevated blood pressure in children). Blood pressure screening would mostly be useful to detect secondary hypertension (eg, due to renal or endocrine disorders) in the very few cases of children with elevated blood pressure who are not obese, considering existing local resources. Drug therapy might be initiated only if blood pressure is persistently elevated despite lifestyle modification. However, unsolved questions regarding drug therapy for elevated blood pressure in children and adolescents concern its absolute cardiovascular disease risk




Figure 3: Prevalence of elevated blood pressure in children and adolescents in Africa stratified by BMI category

Underweight: BMI less than 2 SDs below the WHO growth standard mean for age and sex; normal: BMI between 2 SDs below and 1 SD above the mean; overweight: BMI more than 1 SD above the mean; and obesity: BMI more than 2 SDs above the mean. BMI=body-mass index.

reduction following blood pressure reduction and its effect on prevention of hypertension in adulthood (and thus duration of treatment), as well as risk–benefit ratio.

Results from this study should be interpreted with caution in the context of its limitations. First, definitions of elevated blood pressure differed across studies. Hypertension definition is a major problem in both adult and paediatric studies given the high correlation between blood pressure and growth stature, weight, and sexual maturity in children and adolescents. Hence, the situation might not be well reflected in studies with unacceptable definitions of elevated blood pressure as well as studies involving only participants with high BMI. Even the definition of high blood pressure in children and adolescents considered acceptable in this review might not be suitable for African children and adolescents given that it was derived from the US population.^{11,91} The results from ongoing endeavours to establish international blood pressure references among children and adolescents will help in comparisons of the prevalence of elevated blood pressure in children and adolescents between countries and regions.⁹² Second, we found substantial heterogeneity in estimation of the prevalence of elevated blood pressure across studies.

This heterogeneity might in part be explained by between-study differences in methodology (including setting, device used for blood pressure measurement, and number of measurements) and in population characteristics (eg, proportion of participants with obesity), but it might also reflect true differences of elevated blood pressure burden. Third, we did not have enough data to allow optimal age-stratified analyses. Our results are limited by the wide age band of our study population and the proposed age-stratified analysis. Fourth, UNSD African regions were not uniformly represented, partly owing to difficult retrieval of African medical literature, especially for older articles and those published in local journals.

In conclusion, this study suggests a high prevalence of elevated blood pressure among children and adolescents in Africa, with overweight and obesity being important risk factors. Efforts to address the burden of elevated blood pressure in children and adolescents should mainly focus on primary prevention at the community level by promoting healthy lifestyle and avoiding other cardiovascular risk factors, especially overweight and obesity. More elaborate studies using uniform and reliable diagnostic methods are needed to better estimate

the prevalence of elevated blood pressure and its determinants in children and adolescents within and across countries in Africa.

Contributors

JJN, ME, and JJB conceived the study and, together with JRN, LNA, and AMJ, designed the protocol. JJN, ME, and LNA did the literature search. ME, JJN, and JJB selected the studies and extracted the relevant information. JJB, ME, and JJN synthesised the data. JJN, ME, and JJB wrote the first draft of the paper. JJN, JJB, ME, JRN, LNA, and AMJ critically revised successive drafts of the paper and approved its final version.

Declaration of interests

We declare no competing interests.

References

- WHO. A global brief on hypertension: silent killer, global public health crisis. World Health Day 2013. Geneva: World Health Organization Press, 2013.
- GBD 2015 Risk Factors Collaborators. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 2016; **388**: 1659–724.
- NCD Risk Factor Collaboration. Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population based measurement studies with 19·1 million participants. *Lancet* 2017; **389**: 37–55.
- Santi M, Simonetti BG, Leoni-Foglia CFP, Bianchetti MG, Simonetti GD. Arterial hypertension in children. *Curr Opin Cardiol* 2015; **30**: 403–10.
- Liang YJ, Xi B, Hu YH, et al. Trends in blood pressure and hypertension among Chinese children and adolescents: China Health and Nutrition Surveys 1991–2004. *Blood Press* 2011; **20**: 45–53.
- Peters H, Whincup PH, Cook DG, Law C, Li L. Trends in blood pressure in 9 to 11-year-old children in the United Kingdom 1980–2008: the impact of obesity. *J Hypertens* 2012; **30**: 1708–17.
- Rosner B, Cook NR, Daniels S, Falkner B. Childhood blood pressure trends and risk factors for high blood pressure: the NHANES experience 1988–2008. *Hypertension* 2013; **62**: 247–54.
- McNiece KL, Poffenbarger TS, Turner JL, et al. Prevalence of hypertension and prehypertension among adolescents. *J Pediatr* 2007; **150**: 640–44.
- Bucher BS, Ferrarini A, Weber N, et al. Primary hypertension in childhood. *Curr Hypertens Rep* 2013; **15**: 444–52.
- Redwine KM, Acosta AA, Poffenbarger T, et al. Development of hypertension in adolescents with pre-hypertension. *J Pediatr* 2012; **160**: 98–103.
- Chioloro A, Bovet P, Paradis G. Screening for elevated blood pressure in children and adolescents: a critical appraisal. *JAMA Pediatr* 2013; **167**: 266–73.
- Bao W, Threefoot SA, Srinivasan SR, et al. Essential hypertension predicted by tracking of elevated blood pressure from childhood to adulthood: the Bogalusa Heart Study. *Am J Hypertens* 1995; **8**: 657–65.
- Beckett LA, Rosner B, Roche AF, et al. Serial changes in blood pressure from adolescence into adulthood. *Am J Epidemiol* 1992; **135**: 1166–77.
- Daniels SR, Lloggie JMH, Khoury P, et al. Left ventricular geometry and severe left ventricular hypertrophy in children and adolescents with essential hypertension. *Circulation* 1998; **97**: 1907–11.
- McNiece KL, Gupta-Malhotra M, Samuels J, et al, for the National High Blood Pressure Education Program Working Group. Left ventricular hypertrophy in hypertensive adolescents: analysis of risk by 2004 National High Blood Pressure Education Program Working Group staging criteria. *Hypertension* 2007; **50**: 392–95.
- Raitakari OT, Juonala M, Kahonen M, et al. Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. *JAMA* 2003; **290**: 2277–83.
- Li S, Chen W, Srinivasan SR, et al. Childhood cardiovascular risk factors and carotid vascular changes in adulthood. The Bogalusa Heart Study. *JAMA* 2003; **290**: 2271–76.
- Hartiala O, Magnussen CG, Kajander S, et al. Adolescence risk factors are predictive of coronary artery calcification at middle age: the cardiovascular risk in young Finns study. *J Am Coll Cardiol* 2012; **60**: 1364–70.
- Juonala M, Magnussen CG, Venn A, et al. Influence of age on associations between childhood risk factors and carotid intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study, the Childhood Determinants of Adult Health Study, the Bogalusa Heart Study, and the Muscatine Study for the International Childhood Cardiovascular Cohort (i3C) Consortium. *Circulation* 2010; **122**: 2514–20.
- Hartiala O, Magnussen CG, Kajander S, et al. Adolescence risk factors are predictive of coronary artery calcification at middle age: the cardiovascular risk in young Finns study. *J Am Coll Cardiol* 2012; **60**: 1364–70.
- Litwin M, Niemirska A, Sladowska-Kozłowska J, et al. Regression of target organ damage in children and adolescents with primary hypertension. *Pediatr Nephrol* 2010; **25**: 2489–99.
- Essouma M, Noubiap JJ, Bigna JJ, et al. Hypertension prevalence, incidence and risk factors among children and adolescents in Africa: a systematic review and meta-analysis protocol. *BMJ Open* 2015; **5**: e008472.
- Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ* 2003; **327**: 557–60.
- Barendregt JJ, Doi SA, Lee YY, et al. Meta-analysis of prevalence. *J Epidemiol Community Health* 2013; **67**: 974–78.
- Hoy D, Brooks P, Woolf A, et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol* 2012; **65**: 934–39.
- Cochran WG. The combination of estimates from different experiments. *Biometrics* 1954; **10**: 101–29.
- Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002; **21**: 1539–58.
- Egger M, Davey Smith G, Schneider M, et al. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; **315**: 629–34.
- Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med* 2005; **37**: 360–63.
- Ajite AB, Aladekomo TA, Aderounmu T, Olowu WA. Burden of hypertension and abnormal glomerular permeability in hypertensive school children. *Nephrourol Mon* 2016; **8**: e37568.
- Also U, Asani M, Ibrahim M. Prevalence of elevated blood pressure among primary school children in Kano Metropolis, Nigeria. *Nig J Cardiol* 2016; **13**: 57–61.
- Ansa VO, Anah Mu, Odey FA, et al. Relationship between parental socio-economic status and casual blood pressure in coastal Nigerian adolescents. *West Afr J Med* 2010; **29**: 146–52.
- Aounallah-Skhiri H, El Atti J, Traissac P, et al. Blood pressure and associated factors in a North African adolescent population. A national cross-sectional study in Tunisia. *BMC Public Health* 2012; **12**: 98.
- Awotidebe A, Monyekei MA, Moss SJ, et al. Relationship of adiposity and cardiorespiratory fitness with resting blood pressure of South African adolescents: the PAHL study. *J Hum Hypertens* 2016; **30**: 245–51.
- Ellenga Mbolla BF, Okoko AR, Mabiala Babela JR, et al. Prehypertension and hypertension among schoolchildren in Brazzaville, Congo. *Int J Hypertens* 2014; **2014**: 803690.
- Ghannem H, Trabelsi L, Gaha R, et al. Study of cardiovascular disease risk factors among rural school children in Sousse, Tunisia. *East Mediterr Health J* 2001; **7**: 617–24.
- Harrabi I, Belarbia A, Gaha R, et al. Epidemiology of hypertension among a population of school children in Sousse, Tunisia. *Can J Cardiol* 2006; **22**: 212–16.
- Kidy F, Rutebarika D, Lule SA, et al. Blood pressure in primary school children in Uganda: a cross-sectional survey. *BMC Public Health* 2014; **14**: 1223.
- Kramoh KE, N'goran YN, Aké-Traboulsi E, et al. Prevalence of obesity in school children in Ivory Coast. *Ann Cardiol Angiol (Paris)* 2012; **61**: 145–49.
- Ma C, Kelishadi R, Hong YM, Bovet P, et al. Performance of eleven simplified methods for the identification of elevated blood pressure in children and adolescents. *Hypertension* 2016; **68**: 614–20.

- 41 Monyeki KD, Kemper HCG, Makgae PJ. Relationship between fat patterns, physical fitness and blood pressure of rural South African children: ELLISRAS Longitudinal Growth and Health. *J Hum Hypertens* 2008; **22**: 311–19.
- 42 Motswagole BS, Kruger HS, Faber M, et al. The sensitivity of waist-to-height ratio in identifying children with high blood pressure. *Cardiovasc J Afr* 2011; **22**: 208–11.
- 43 Odetunde OI, Neboh EE, Chinawa JM, et al. Elevated arterial blood pressure and body mass index among Nigerian preschool children population. *BMC Pediatrics* 2014; **14**: 64.
- 44 Odey F, Anah M, Ansa V, Ogbeche J, Meremikwu M, Ekanem E. Pre-hypertension and hypertension in apparently healthy adolescents in Calabar, Nigeria. *Glob J Community Med* 2009; **2**: 13–20.
- 45 Okoh BA, Alikor EA, Akani N. Prevalence of hypertension in primary school-children in Port Harcourt, Nigeria. *Paediatr Int Child Health* 2012; **32**: 208–12.
- 46 Okpere AN, Anochie IC, Eke FU. Pattern of blood pressure and hypertension in adolescents in Port Harcourt, Nigeria. *West Afr J Med* 2013; **32**: 93–98.
- 47 Oyewole OO, Oritogun KS. Pre-hypertension and hypertension in adolescence: how much does it occur in a Nigerian community? *West Afr J Med* 2012; **31**: 71–75.
- 48 Sadoh WE, Omuemu VO, Sadoh AE, et al. Blood pressure percentiles in a group of Nigerian school age children. *Niger J Paed* 2014; **41**: 223–28.
- 49 Ujunwa FA, Ikefuna AN, Nwokocha ARC, et al. Hypertension and prehypertension among adolescents in secondary schools in Enugu, South East Nigeria. *Ital J Pediatr* 2013; **39**: 70.
- 50 Xi B, Bovet P, Hong YM, et al. Recent blood pressure trends in adolescents from China, Korea, Seychelles and the United States of America, 1977–2012. *J Hypertens* 2016; **34**: 1948–58.
- 51 Aboul Ella NA, Shehab DI, Ismail MA, et al. Prevalence of metabolic syndrome and insulin resistance among Egyptian adolescents 10 to 18 years of age. *J Clin Lipidol* 2010; **4**: 185–95.
- 52 Akingugbe FM, Akinwolere AOA, Kayode CM. Blood pressure patterns in Nigerian adolescents. *West Afr J Med* 1999; **18**: 196–202.
- 53 Akor F, Okolo SN, Okolo AA. Blood pressure and anthropometric measurements in healthy primary school entrants in Jos, Nigeria. *S Afr J Child Health* 2010; **4**: 42–45.
- 54 Benmohammed K, Nguyen MT, Khensal S, et al. Arterial hypertension in overweight and obese Algerian adolescents: role of abdominal obesity. *Diabetes Metab* 2011; **37**: 291–97.
- 55 Chiolerio A, Madeleine G, Gabriel A, et al. Prevalence of elevated blood pressure and association with overweight in children of a rapidly developing country. *J Hum Hypertens* 2007; **21**: 120–27.
- 56 Ejike CE, Ugwu CE, Ezeanyika LU. Variations in the prevalence of point (pre)hypertension in a Nigerian school-going adolescent population living in a semi-urban and an urban area. *BMC Pediatrics* 2010; **10**: 13.
- 57 Ghannem H, Ahmed BA, Khellifa L, et al. Phénomène de “tracking” des facteurs de risque cardiovasculaires chez l’enfant en milieu scolaire étude de cohorte sur 4 ans à Sousse en Tunisie. *Tunis Med* 2005; **83**: 404–08.
- 58 Kagura J, Adair LS, Musa MG, et al. Blood pressure tracking in urban black South African children: birth to twenty cohort. *BMC Pediatrics* 2015; **15**: 78.
- 59 Kane A, Ly M, Sarr M, et al. Etude de la pression artérielle et de l’index de masse corporelle chez l’enfant et l’adolescent en milieu rural à Thiadiaye (Sénégal). *Dakar Med* 1998; **43**: 83–89.
- 60 Ladapo TA, Fajolu IB, Adeniyi OF, et al. Blood pressure to height ratio as a screening tool for prehypertension and hypertension in adolescents. *Niger J Clin Pract* 2016; **19**: 401–06.
- 61 Longo-Mbenza B, Ngiyulu R, Bayekula M, et al. Low birth weight and risk of hypertension in African school children. *J Cardiovasc Risk* 1999; **6**: 311–14.
- 62 Mehdad S, Hamrani A, Kari KE, et al. Prevalence of elevated blood pressure and its relationship with fat mass, body mass index and waist circumference among a group of Moroccan overweight adolescents. *Obes Res Clin Pract* 2013; **7**: e284–89.
- 63 Mijinyawa M, Llyiasu Z, Borodo MM. Prevalence of hypertension among teenage students in Kano, Nigeria. *Niger J Med* 2008; **17**: 173–78.
- 64 Nkeh-Chungag BN, Sekokotla AM, Sewani-Rusike C, et al. Prevalence of hypertension and pre-hypertension in 13-17 year old adolescents living in Mthatha—South Africa: a cross-sectional study. *Cent Eur J Public Health* 2015; **23**: 59–64.
- 65 Musa DI, Williams GA. Cardiorespiratory fitness, fatness, and blood pressure associations in Nigerian youth. *Med Sci Sports Exer* 2012; **44**: 1978–85.
- 66 Owuoye OB, Olawale OA, Tella BA, Ajuluchukwu JN, Akinbo SR. Prevalence of hypertension and pre-hypertension in male adolescent football: a cross-sectional cohort study of Nigerian players. *Am J Hypertens Res* 2013; **1**: 26–28.
- 67 Odunaiya NA, Louw QA, Grimmer KA. Are lifestyle cardiovascular disease risk factors associated with pre-hypertension in 15–18 years rural Nigerian youth? A cross-sectional study. *BMC Cardiovasc Disord* 2015; **15**: 144.
- 68 Oduwale AA, Ladapo TA, Fajolu IB, et al. Obesity and elevated blood pressure among adolescents in Lagos, Nigeria: a cross-sectional study. *BMC Public Health* 2012; **12**: 616.
- 69 Okagua J, Anochie FC. Blood pressure profile and hypertension in adolescents in Port Harcourt, Southern Nigeria. *Afr J Paed Nephrol* 2014; **1**: 77–82.
- 70 Okagua J, Akani N. Prevalence of hypertension in school going adolescents in rural areas of Rivers State, south-South Nigeria. *Nigerian Health J* 2014; **14**: 157–64.
- 71 Omisore B, Omisore AG, Abioye-Kuteyi EA. Gender comparisons of adolescents’ anthropometry and blood pressure in Osun State, South-Western Nigeria. *Int J Adolesc Med Health* 2015; **27**: 247–51.
- 72 Oyeyemi AY, Usman MA, Oyeyemi AL, et al. Casual blood pressure of adolescents attending public secondary schools in Maiduguri, Nigeria. *Clin Hypertens* 2015; **21**: 16.
- 73 Salvi P, Meriem C, Temmar M, et al. Association of current weight and birth weight with blood pressure levels in Saharan and European teenager populations. *Am J Hypertens* 2010; **23**: 379–86.
- 74 Silva AB, Capingana DP, Magalhães P, et al. Cardiovascular risk factors in pre-pubertal schoolchildren in Angola. *Cardiovasc J Afr* 2016; **27**: 315–21.
- 75 Steyn K, de Wet T, Richter L, et al. Cardiovascular disease risk factors in 5-year-old urban South African children—the birth to ten study. *S Afr Med J* 2000; **90**: 719–26.
- 76 Uwaezuoke SN, Okoli CV, Ubesie AC, et al. Primary hypertension among a population of Nigerian secondary school adolescents: prevalence and correlation with anthropometric indices: a cross-sectional study. *J Clin Pract* 2016; **19**: 649–59.
- 77 de Moraes AC, Lacerda MB, Moreno LA, Horta BL, Carvalho HB. Prevalence of high blood pressure in 122,053 adolescents: a systematic review and meta-regression. *Medicine (Baltimore)* 2014; **93**: e232.
- 78 Ataklte F, Erqou S, Kaptoge S, Taye B, Echouffo-Tcheugui JB, Kengne AP. Burden of undiagnosed hypertension in Sub-Saharan Africa: a systematic review and meta-analysis. *Hypertension* 2015; **65**: 291–98.
- 79 Alwan A, Armstrong T, Bettcher D, et al. Global status report on noncommunicable diseases 2010. Geneva: World Health Organization, 2011.
- 80 Muthuri SK, Francis CE, Wachira LJ, et al. Evidence of an overweight/obesity transition among school-aged children and youth in Sub-Saharan Africa: a systematic review. *PLoS One* 2014; **9**: e92846.
- 81 Dinsa GD, Goryakin Y, Fumagalli E, Suhrcke M. Obesity and socioeconomic status in developing countries: a systematic review. *Obes Rev* 2012; **13**: 1067–79.
- 82 Pisa PT, Pisa NM. Economic growth and obesity in South African adults: an ecological analysis between 1994 and 2014. *Eur J Public Health* 2017; **27**: 404–09.
- 83 Witoszynska B, Krzyzaniak A, Krzyzaniak A, Kaczmarek M, Siwinska A. Environmental variation in the prevalence of hypertension in children and adolescents—is blood pressure higher in children and adolescents living in rural areas. *Ann Agric Environ Med* 2017; **24**: 129–33.
- 84 Aounallah-Skhiri H, Romdhane H, Traissac P, et al. Nutritional status of Tunisian adolescents: associated gender, environmental and socioeconomic factors. *Public Health Nutr* 2008; **11**: 1306–17.
- 85 Bareto SM, Miranda JJ, Figueroa JP, et al. Epidemiology in Latin America and the Caribbean: current situation and challenges. *Int J Epidemiol* 2012; **41**: 557–71.

- 86 Shin JH, Lee G, Kim JS, et al. Effects of the 2008 global economic crisis on national health indicators: results from the Korean National Health and Nutrition Examination Survey. *Korean J Fam Med* 2015; **36**: 162–67.
- 87 Leng B, Jin Y, Li G, Chen L, Jin N. Socioeconomic status and hypertension: a meta-analysis. *J Hypertens* 2015; **33**: 221–29.
- 88 Kagura J, Adair LS, Pisa PT, Griffiths PL, Pettifor JM, Norris SA. Association of socioeconomic status change between infancy and adolescence, and blood pressure, in South African young adults: birth to twenty cohort. *BMJ Open* 2016; **6**: e008805.
- 89 Bruner MW, Lawson J, Pickett W, Boyce W, Janssen I. Rural Canadian adolescents are more likely to be obese compared with urban adolescents. *Int J Pediatr Obes* 2008; **3**: 205–11.
- 90 Sjöberg A, Moraes L, Yngve A, Poortvliet E, Al-Ansari U, Lissner L. Overweight and obesity in a representative sample of schoolchildren—exploring the urban–rural gradient in Sweden. *Obes Rev* 2011; **12**: 305–14.
- 91 National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* 2004; **114**: 555–76.
- 92 Xi B, Zong X, Kelishadi R, et al. Establishing international blood pressure references among nonoverweight children and adolescents aged 6 to 17 years. *Circulation* 2016; **133**: 398–408.